

## PRE-PACKHOUSE DRENCH

### General recommendation: Thiabendazole + Pyrimethanil + 2,4-D + Peracetic acid

1. The aim is to protect fruit from decay during storage and degreening.
2. Thiabendazole and pyrimethanil may be replaced with other fungicides provided they will not be applied again in the packhouse.
3. NB: ENSURE THAT FUNGICIDES FROM DIFFERENT FRAC GROUPS ARE SELECTED.

### Mixing protocol

1. Fill water tank to 80% capacity and start pump(s).
2. Pre-suspend fungicides individually in lukewarm water (35 – 40°C).
3. Order of mixing: SG – SC – EC – SL
4. Fill water tanks to full capacity.
5. Let the system run for 15 min before commencing treatment to facilitate thorough mixing.

### Operation

1. The drench system must deliver at least 500 l drench mixture/bin/minute.
2. Exposure time must be 1 to 3 minutes.
3. Replace mixture after 150 bins (1 000 l), or 200 bins (2 000 l), or 300 bins (3 000 l).
4. Mixtures can be sanitised with a registered and compatible sanitiser.
5. Constant agitation of the mixture is necessary.

## VOOR-PAKHUISSTORT

### Algemene aanbeveling: Thiabendasool + Pirimetaniel + 2,4-D + Perasynsuur

1. Die doel is om vrugte te beskerm teen bederf tydens opberging en ontgroening.
2. Thiabendasool en Pirimetaniel kan vervang word deur ander swamdoders, maar dan mag die swamdoders nie weer in die pakhuis toegedien word nie.
3. NB: VERSEKER DAT SWAMDODERS VAN VERSKILLENDE FRAC GROEPE GESELEKTEER WORD.

### Mengprotokol

1. Vul tenk tot 80% van kapasiteit en skakel pomp(e) aan.
2. Suspendeer individuele fungisiedes vooraf in lou water (35 – 40°C).
3. Volgorde van meng: SG – SC – EC – SL
4. Vul tenk tot volle kapasiteit.
5. Laat die sisteem sirkuleer vir 15 minute voor behandeling begin ten einde deeglike vermenging te verseker.

### Operasioneel

1. Die sisteem moet minstens 500 l stortmengsel/massakrat ("bin")/minuut lewer.
2. Blootstellingstyd moet 1 tot 3 min wees.
3. Vervang mengsel na 150 kratte (1 000 l), of 200 kratte (2 000 l), of 300 kratte (3 000 l).
4. Mengsels kan gesanteer word deur 'n geregistreerde en verenigbare saniteerder te gebruik.
5. Dit is noodsaaklik dat die mengsel voortdurend geroer word.

For further specifications contact the suppliers of the above mentioned products.

**ALWAYS REFER TO PRODUCT LABELS FOR APPLICATION DETAILS.**

MRL list and restrictions: Visit [www.cga.co.za](http://www.cga.co.za) or email [ph@cga.co.za](mailto:ph@cga.co.za) for "RECOMMENDED USAGE RESTRICTIONS FOR PLANT PROTECTION PRODUCTS ON SOUTHERN AFRICAN EXPORT CITRUS"