

CITRUS BIOSECURITY THREAT: Yellow Vein Clearing Disease



Symptoms

Symptoms are most prominent on lemons, with foliar symptoms of vein clearing and leaf crinkling, with irregular, elongated chlorotic spots, predominantly visible on young growth. Veins on the underside of the leaf become water-soaked and turn brown. Foliar symptoms are accompanied by leaf drop.

On other citrus types, irregular ringspots may be seen on leaves, and mosaic-like patterns on fruit.

In severe infections, trees die back and fruit are malformed, causing reduced fruit quality.

Symptoms are more visible in the cooler periods of spring and autumn and less noticeable in hot environments. Yellow vein clearing disease can be asymptomatic on some citrus cultivars.



Etrog citron (*C. medica*)



Pineapple (*C. sinensis*)



Kinnow mandarin (*C. reticulata*)



Meena *et al.*, Biotech (2019)

Malta (*C. sinensis*)

Vein clearing and flecking on 'Etrog' citron and atypical chlorotic ringspots on 'Pineapple' sweet orange, 'Malta' sweet orange and 'Kinnow' mandarin, similar to symptoms associated with ICRSV

The disease

- Yellow vein clearing disease is caused by citrus yellow vein clearing virus (CYVCV).
- The disease is associated with poor tree vigour, reduced yields and decreased marketability of fruit.
- There is no treatment once trees become infected. Infected trees need to be eradicated



Vein clearing and flecking on 'Eureka' lemon



Possible confusion with

Atypical chlorotic ringspot, similar to those induced by Indian chlorotic ringspot virus (ICRSV), were recorded on 'Kinnow' mandarin and 'Malta' sweet orange.

[Defoliation of CYVCV infected tree >](#)



Host range


- All *Citrus* spp.
- Severe symptoms are observed on 'Eureka' lemon and sour orange
- Grapevines and a range of herbaceous plants and weed species

Current distribution

- Turkey
- China
- India
- Iran
- Pakistan
- California, U.S.A.


Method of spread

- **Infected plant propagation material**
Citrus propagation material (trees, cuttings, grafts, budwood, rootstock seedlings) and propagation material of other hosts
- **Viruliferous insect vectors**
Aphids: *Aphis craccivora* on non-citrus hosts, *Aphis spiraecola* on citrus and non-citrus hosts
Whitefly: *Dialeurodes citri*
- **NOT fruit or seed transmitted**



CABI

Aphis craccivora



CABI

Aphis spiraecola



CABI

Dialeurodes citri

Preventative actions

- Quarantine procedures for importation of citrus propagation material and other hosts
- Plant certified disease-free citrus trees
- Awareness and surveillance to ensure early detection and rapid implementation of control measures
- Do not bring illegal plant material into South Africa and onto your farm

For more information on this disease, or if you find anything unusual, contact Wayne Kirkman from CRI's Biosecurity Division: waynek@cri.co.za, 084 458 0349

Disclaimer: The material in this publication is for general information only, and no person should act, or fail to act on the basis of this material without obtaining professional advice. Citrus Research International (CRI) and all persons acting for CRI expressly disclaim liability with respect to anything done in reliance on this publication.