

RECOMMENDED USAGE RESTRICTIONS FOR PLANT PROTECTION PRODUCTS ON SOUTHERN AFRICAN EXPORT CITRUS

Compiled by:

Paul Hardman, Citrus Growers Association of southern Africa, tel 031-7652514, ph@cga.co.za,
Paula Bester, Citrus Growers Association of southern Africa, tel 031-7652514, pb@cga.co.za,
and Vaughan Hattingh, Citrus Research International, tel 021-8828553, vh@cri.co.za
December 2023 (Number 56)

The purpose of these restrictions is to ensure compliance with residue tolerances in the countries to which southern African citrus is exported. The recommendations on container labels are based on the **current registrations** of plant protection products. In terms of the SA Act 36 of 1947 (and equivalent legislation in neighbouring countries) the registration requirements must be adhered to at all times. However, the requirements of importing countries have made it necessary to introduce **further restrictions** in order to comply with maximum residue limits (MRLs).

The active ingredients of the products are listed alphabetically together with an illustrative brand name. In the case of commodity products which may be sold under different brand names, only a single name is shown for convenience. This does not imply endorsement of the particular brand concerned.

Due to the multiple fruit set pattern of lemons, plant protection products may only be used as specified during the first spring **blossoming and fruit set** period. Special caution should be exercised to ensure that pesticides applied to protect later sets do not result in unacceptable residues in fruit remaining on the trees from earlier sets.

The following general statements are applicable to the use of plant protection products on all cultivars:

- * Growers should ensure that these restrictions are kept handy and are consulted before the application of plant protection products.
- * The application in accordance with current label requirements will in many instances not ensure that export requirements will be met. The restrictions specified are applicable in addition to the label requirements.
- * The withholding periods specified on product labels provide an indication of the ability of treatments to conform to South African residue levels. Since overseas requirements are generally more stringent and these withholding periods are not adequate unless specifically mentioned in this document.
- * These restrictions apply to the period during which fruit is present on trees (between blossom and harvest) and not to the period between harvest and the onset of blossom, during which time standard label requirements apply.
- * All usages apply to normal blossom situations. Under conditions where blossoming occurs over an extended period, a more conservative approach must be adopted. Under these conditions treatments should be timed according to the early portion of the blossom.
- * Particular attention must be given to ensuring that spray machinery is calibrated to apply the correct spray volumes in relation to tree size and that spray operators are trained in the handling and application of plant protection products.
- * All treatments referred to above must be applied at the registered concentrations.
- * Alternation of products, where applicable with reference to the restrictions, will reduce the risk of excessive residues of any one chemical and will also reduce selection pressure for resistance.
- * The additional restrictions in this document do not necessarily provide an indication of the compatibility of the products with integrated pest management and good agricultural practice.
- * The addition of oil to a treatment, if not registered as such, should be avoided as this may increase the residue level.

All exporting growers should keep accurate spray records so that in the event of exceeding MRLs the reasons can be determined. These records should be retained in safe-keeping for at least 3 years.

Growers are strongly urged to abide by these restrictions to minimise the risk of residue tolerances being exceeded. However, it must be noted that no absolute guarantee can be given that even by following these guidelines export residue tolerances will in all instances not be exceeded. The efficacy and integrated pest management compatibility of plant protection products listed here are additional considerations that users should bear in mind and which are not covered in this document whatsoever. This document has been compiled with information presently available and in good faith, but with the express condition that the authors, Citrus Research International and Citrus Growers' Association of Southern Africa, accept no responsibility whatsoever for any loss or damage resulting directly or indirectly from the use thereof.

SUMMARY TABLE OF RECOMMENDED USAGE RESTRICTIONS

PRODUCT	All markets (including EU) except where other restrictions are specified	CODEX (A) ^a	CODEX (B) ^b	CANADA	U S A	JAPAN	Other ^d
Acephate/ Protectostem	70d PHI	-	40d PHI as registered.	-	-	-	South Korea: 70d PHI for lemons, oranges and grapefruit and 40d PHI for soft citrus.
Acetamiprid/ Mospilan	150d PHI as registered.	-	-	-	-	-	South Korea: 150d PHI as registered.
Acrinathrin/ Rufast	Not later than 90% petal fall.	-	-	-	-	-	-
Amitraz/ Mitac	Not later than 90% petal fall.	-	-	150d PHI	-	28d PHI as registered.	-
Avermectin/ Agrimec	7d PHI as registered.	-	-	-	-	-	-
Azadirachtin	See Pyrethrins	-	-	-	-	-	-
Azinphos-methyl/ Gusathion	Not later than 90% petal fall.	120d PHI	120d PHI	21d PHI as registered.	-	21d PHI as registered.	GSO: 120d PHI.
Azoxystrobin/ Ortiva	Pre-harvest: 77d PHI as registered on grapefruit, oranges and soft citrus, and 14d as registered on lemons. Post-harvest: Post-harvest as registered.	-	-	-	-	-	China: Pre-harvest: 77d PHI as registered on grapefruit, oranges and soft citrus, and 14d as registered on lemons. Post-harvest: Not permitted.
<i>Bacillus thuringiensis</i> / Dipel	0d PHI as registered.	-	-	-	-	-	-
<i>Beauveria bassiana</i> / BroadBand	0d PHI as registered.	-	-	-	-	-	-
Bromopropylate/ Acarol	Not later than 90% petal fall.	-	-	21d PHI	-	21d PHI	GSO: 21d PHI.
Buprofezin/ Applaud	Not later than mid- October for oranges and grapefruit and not later than 90% petal fall for lemons and soft citrus.	45d PHI as registered.	45d PHI as registered.	45d PHI as registered.	45d PHI as registered.	45d PHI as registered.	45d PHI as registered. South Korea: Not later than mid- October for grapefruit and 45d PHI as registered for other citrus.
Cadusaphos/ Rugby	0d PHI as registered.	-	-	-	-	-	-

PRODUCT	All markets (including EU) except where other restrictions are specified	CODEX (A) ^a	CODEX (B) ^b	CANADA	U S A	JAPAN	Other ^d
Carbendazim (Bavistin, Bendazid, Knowin, Benomyl, Spotless)	120d PHI for oranges and grapefruit and 90d PHI for other citrus. (Not later than 90% petal fall for oranges and grapefruit and 90d PHI for lemons and mandarins) ^{EU5}	90d PHI for oranges and not later than 90% petal fall for other citrus.	14d PHI as registered.	14d PHI as registered.	Not later than 90% petal fall.	14d PHI as registered.	South Korea: Not later than 90% petal fall for oranges and grapefruit and 120d PHI as registered for lemons and 14d as registered for soft citrus. GSO: 90d PHI for oranges and 14d PHI as registered for other citrus. Thailand: 120d PHI for oranges and not later than 90% petal fall for other citrus. China: 14d PHI as registered for oranges and soft citrus and 90d PHI for lemons and grapefruit. Bangladesh: 90d PHI for oranges and not later than 90% petal fall for other citrus.
Chlorfenapyr/ Hunter	Medium cover spray: Before calyx closure (\pm 3 weeks after petal fall) as registered. Bait spray application (30ml): Not later than mid-December and a 140d PHI as registered.	-	-	-	-	-	-
Chlorpyrifos/ Dursban	Not later than 90% petal fall for sprays, 0d PHI for soil and stem applications as registered. ^f	-	-	60d PHI for sprays as registered, 0d PHI for soil and stem applications as registered.	-	60d PHI for sprays as registered, 0d PHI for soil and stem applications as registered.	China, GSO, India, South Korea, Switzerland, Singapore, Vietnam: 60d PHI for sprays as registered, 0d PHI for soil and stem applications as registered. Thailand: Not later than 90% petal fall.
Chlorantraniliprole (Rynaxypyr)/ Coragen	7d PHI as registered.	-	-	-	-	7d PHI as registered. Not later than 90% petal fall for soft citrus.	Norway: see note g.
Clothianidin/ Dantop	120d PHI as registered	-	-	-	-	-	-
Copper	14d PHI as registered	-	-	-	-	-	-
Cyantraniliprole / Exirel	7d PHI as registered for foliar applications and 1d PHI as registered for bait sprays.	-	-	-	-	-	Taiwan: Not later than 90% petal fall.

PRODUCT	All markets (including EU) except where other restrictions are specified	CODEX (A) ^a	CODEX (B) ^b	CANADA	U S A	JAPAN	Other ^d
Cyhexatin (Azocyclotin)/ Sipcatin	Only to be used after harvest and before the onset of blossom.	-	-	-	Not permitted	-	Taiwan and Thailand: Not permitted.
Cypermethrin	28d PHI as registered. (Not later than 90% petal fall for oranges and 28d PHI as registered for other citrus). ^{EU1}	-	-	-	28d PHI as registered for alpha and zeta-Cypermethrin formulations. Not later than 90% petal fall for Cypermethrin ^c	-	-
Dichlorprop-p/ Corasil P	90d PHI as registered.	-	-	-	-	-	
Dichlorvos/ Devipan	21d PHI	-	-	-	-	-	-
Dicofol/ Kelthane	Not later than 90% petal fall.	-	14d PHI. Russia: Not later than 90% petal fall.	-	-	14d PHI	
Difenoconazole/ Score	Not later than 90% petal fall as registered.	-	-	-	-	-	-
Dimethoate/ Rogor	Not later than 50% petal fall for foliar applications and not later than white bud stage for soil applications.	14d PHI for foliar applications and 42d PHI for soil applications as registered for mandarins. For other citrus not later than 50% petal fall for foliar applications and not later than white bud stage for soil applications.	-	14d PHI for foliar applications and 42d PHI for soil applications as registered.	14d PHI for foliar applications and 42d PHI for soil applications as registered.	14d PHI for foliar applications and 42d PHI for soil applications as registered.	GSO: 14d PHI for foliar applications and 42d PHI for soil applications as registered.

PRODUCT	All markets (including EU) except where other restrictions are specified	CODEX (A) ^a	CODEX (B) ^b	CANADA	U S A	JAPAN	Other ^d
Dimethyl Didecylammonium Chloride/ Sporekill	Post-harvest: not permitted. Pre-harvest: 160d PHI.	-	Pre-harvest and post-harvest as registered.	-	-	Post-harvest: not permitted. Pre-harvest: 160d PHI ^k	-
Dithiocarbamates (Mancozeb/Maneb)	21d PHI or 28d PHI as registered.	Not later than 90% petal fall for Lemons and Grapefruit and 21d or 28d PHI as registered for Soft Citrus and Oranges.	-	Not later than end December and only where packhouses have either a non-recycling high pressure spray or non-recycling Deccosol foam curtain and regular (twice daily) cleaning of brushes.	Not later than 90% petal fall for Oranges, Lemons and Grapefruit and 21d or 28d PHI as registered for Soft Citrus.	Not later than end-January.	South Korea: 21d PHI or 28d PHI as registered for soft citrus, and lemons. Not later than end December and only where packhouses have either a non-recycling high pressure spray or non-recycling Deccosol foam curtain and regular (twice daily) cleaning of brushes for grapefruit and oranges. India: Not later than end December and only where packhouses have either a non-recycling high pressure spray or non-recycling Deccosol foam curtain and regular (twice daily) cleaning of brushes. Thailand: 21d PHI or 28d PHI as registered. Bangladesh: Not later than 90% petal fall. GSO countries: Not later than 90% petal fall for Lemons and Grapefruit and 21d or 28d PHI as registered for Soft Citrus and Oranges.
Dipotassium phosphate	7d PHI as registered	-	-	-	-	-	-
(E)-8-Dodecen-1-yl acetate + (Z)-8-Dodecen-1-yl acetate/ Checkmate	0d PHI as registered.	-	-	-	-	-	-
Emamectin Benzoate/ Warlock	14d PHI as registered.	-	-	-	-	-	-
Ethephon/ Ethrel	Not permitted	-	As registered	As registered	-	Post-harvest use not permitted.	-

PRODUCT	All markets (including EU) except where other restrictions are specified	CODEX (A) ^a	CODEX (B) ^b	CANADA	U S A	JAPAN	Other ^d
Ethoprophos/ MOCAP	0d PHI as registered.	-	-	-	-	-	-
Etoazole/ Smite	28d PHI as registered. (Not later than 90% petal fall) ^{EU4}	-	-	-	Not later than 90% petal fall for lemons and grapefruit and 28d PHI as registered for oranges and soft citrus.	-	South Korea: Not later than 90% petal fall for oranges and grapefruit and 28d PHI as registered for lemons and soft citrus.
Fenamiphos/ Nemacur	150d PHI as registered.	-	-	-	-	-	-
Fenazaquin	56d PHI as registered.	-	-	-	-	-	-
Fenbutatin-oxide/ Torque	Not later than 90% petal fall.	7d PHI as registered.	7d PHI as registered.	7d PHI as registered.	7d PHI as registered.	7d PHI as registered.	7d PHI as registered. South Korea: Not later than 90% petal fall.
Fenpropathrin/ Meothrin	28d PHI as registered (180d PHI) ^{EU2}	-	-	-	-	-	-
Fenpropathrin + Phenthoate/ Meothrin + Elsan	Not later than 90% petal fall.	-	-	-	-	-	-
Fenpyroximate/ Mitigate	28d PHI as registered.	28d PHI as registered for lemons and grapefruit and not later than 90% petal fall for oranges and mandarins	-	-	-	-	-
Fipronil/ Regent	Before calyx closure (± 3 weeks after petal fall) as registered.	-	-	-	-	-	-
Fluopyram / Velum	120d PHI as registered.	-	-	-	-	-	-
Flutriafol/ Impact	Not later than 90% petal fall as registered.	Not later than 90% petal fall.	-	-	Not later than 90% petal fall.	Not later than 90% petal fall.	South Korea: Not later than 90% petal fall.
Formetanate/ Dicarzol	90d PHI as registered.	-	-	-	-	-	-
Fosetyl-Al/Aliette	25d or 0d PHI as registered.	-	-	-	-	-	-
Fludioxonil/ Teacher	Post-harvest as registered.	-	-	-	-	-	Taiwan: T1
Fosthiazate/ Nemathorin	43d PHI as registered.	-	-	-	-	-	-
Furfural/ Crop Guard	43d PHI as registered.	-	-	-	-	-	-

PRODUCT	All markets (including EU) except where other restrictions are specified	CODEX (A) ^a	CODEX (B) ^b	CANADA	U S A	JAPAN	Other ^d
Gibberellic Acid	15d PHI as registered.	-	-	-	-	-	-
Granulovirus (Cryptogran, Cryptex)	0d PHI as registered.	-	-	-	-	-	-
Guazatine	Not Permitted	Post-harvest as registered.	Post-harvest as registered.	-	-	-	GSO: Post-harvest as registered.
Helicoverpa armigera nucleopolyhedrovirus/ Helicovir	0d PHI as registered.	-	-	-	-	-	-
Imazalil (Chloramizol)	Post-harvest as registered.	-	-	-	-	-	-
Imidacloprid/ Confidor	212d PHI as registered (Not to be used on bearing trees for EU fruit in the market from July 2022). ^{EU3}	212d PHI as registered.	212d PHI as registered.	212d PHI as registered.	212d PHI as registered.	212d PHI as registered.	Great Britain ^m : 212d PHI as registered. Taiwan: 212d PHI as registered for oranges and not later than 90% petal fall for other citrus. Thailand: 212d PHI as registered.
Iprodione/ Rovral (Dicarboxamil)	Not later than 90% petal fall.	-	115d PHI as registered.	-	-	-	GSO, Switzerland and Taiwan: Not later than 90% petal fall. South Korea: 115d PHI as registered for soft citrus and lemons and not later than 90% petal fall for other citrus.
Isazophos/ Miral	56d PHI as registered.	-	-	-	-	-	-
Kresoxim-methyl/ Strobby	56d PHI for grapefruit & oranges. Not later than 90% petal fall for lemons & soft citrus.	-	56d PHI	56d PHI for soft citrus, and not later than 90% petal fall for other citrus.	Not later than 90% petal fall.	56d PHI	Not later than 90% petal fall. South Korea: 115d PHI as registered for soft citrus and not later than 90% petal fall for other citrus.
Mercaptothion/ Malathion (fruit fly baiting only)	Only dilute concentration, do not use higher concentrations. 7d PHI.	-	-	14d PHI	-	-	South Korea: 7d PHI for oranges and 28d PHI for other citrus. Thailand: 14d PHI for grapefruit and 7d PHI for other citrus. Switzerland: 28d PHI.
Metalaxyl M/ Ridomil Gold	30d PHI as registered (Not later than 90% petal fall for lemons) ^{EU6}	-	-	-	-	-	South Korea: Not later than 90% petal fall.
Methamidophos/ Citrimet	60d PHI	-	21d PHI as registered.	21d PHI as registered.	-	-	-

PRODUCT	All markets (including EU) except where other restrictions are specified	CODEX (A) ^a	CODEX (B) ^b	CANADA	U S A	JAPAN	Other ^d
Methidathion/ Ultracide	Not later than 90% petal fall.	-	56d PHI as registered.	56d PHI as registered.	-	56d PHI as registered.	GSO: 56d PHI as registered.
Methiocarb/ Mesurol	Not later than the end of January.	-	-	-	-	-	South Korea: 21d PHI as registered for soft citrus and not later than the end of January for others. Taiwan: Not later than 90% petal fall. GSO: Not later than the end of January.
Methomyl (Thiodicarb) / Lannate	60d PHI for all registered usages.	28d PHI for all registered usages.	28d PHI for all registered usages.	28d PHI for all registered usages.	28d PHI for all registered usages.	28d PHI for all registered usages.	GSO: 28d PHI for all registered usages.
Methoxyfenozide / Runner	30d PHI as registered.	-	-	-	-	-	-
Methyl-parathion / Pennacap	Not later than 50% petal fall.	-	-	-	-	-	-
Mevinphos	28d PHI	-	3d PHI as registered.	3d PHI as registered .	-	-	South Korea: 3d PHI as registered.
Orange Oil/ Prev-Am	0d PHI as registered.	-	-	-	-	-	-
Omethoate/ Folimat	Apply no more than once in a season, not later than beginning of December and ensure at least a 150d PHI.	-	-	-	-	-	-
Parathion/ Parathion	Not later than 50% petal fall.	-	-	-	-	Not later than 4 weeks after petal fall.	-
Permethrin/ Last Call	0d PHI as registered.	-	-	-	-	-	-
Phenthoate/ Elsan	Not later than 50% petal fall.	-	-	-	-	-	-
Phosphorous acid	0d PHI, 14d PHI or 28d PHI as registered.	-	-	-	-	-	-
Pirimicarb/ Aphox	14d PHI as registered.	-	-	Not later than 90% petal fall.	Not later than 90% petal fall.	14d PHI as registered for oranges and not later than 90% petal fall for other citrus.	Switzerland, Singapore, South Korea, Vietnam: Not later than 90% petal fall.

PRODUCT	All markets (including EU) except where other restrictions are specified	CODEX (A) ^a	CODEX (B) ^b	CANADA	U S A	JAPAN	Other ^d
Potassium bicarbonate/ Agri-Cure SP	0d PHI as registered.	-	-	-	-	-	-
Prochloraz	Not permitted.	Post-harvest as registered.	Post-harvest as registered.	-	-	-	-
Profenofos/ Selecron	Between blossom and harvest, use Selecron only once and not more than 100mℓ/100ℓ water at not later than 50% petal fall ^Y	-	-	-	-	-	-
Propiconazole / Propicure	Not permitted.	Post-harvest as registered.	Post-harvest as registered.	Post-harvest as registered.	Post-harvest as registered.	Post-harvest as registered.	Not permitted for Hong Kong, India, Vietnam, GSO countries. Taiwan: Post-harvest as registered for oranges, lemons and grapefruit and not permitted on soft citrus.
Prothiofos/ Tokuthion	Between blossom and harvest, Prothiofos should be used only once and not later than 90% petal fall.	-	-	-	-	-	-
Pymetrozine/ Chess	Not later than 90% petal fall.	-	-	42d PHI as registered for Oranges, Soft Citrus and Lemons, and not later than 90% petal for Grapefruit.	-	-	-
Pyraclostrobin/ Cabrio	45d PHI as registered.	-	-	-	-	-	Taiwan: Not later than 90% petal fall for soft citrus and 45d PHI as registered for other citrus.
Pyrethrins (incl natural Pyrethrum) / Erador	2d PHI as registered.	Not later than 90% petal fall.	Not later than 90% petal fall.	2d PHI as registered for oranges and not later than 90% petal fall for other citrus.	2d PHI as registered for oranges and not later than 90% petal fall for other citrus.	-	China, Hong Kong, Russia, South Korea, Taiwan, Thailand and GSO countries: Not later than 90% petal fall.

PRODUCT	All markets (including EU) except where other restrictions are specified	CODEX (A) ^a	CODEX (B) ^b	CANADA	U S A	JAPAN	Other ^d
Pyrimethanil / Philabuster	Pre-harvest: 90d PHI as registered. Post-harvest as registered. Do not use in HOT water bath.	Pre-harvest: 90d PHI as registered. Post-harvest as registered. Do not use in HOT water bath and not recommended for wax application.	-	-	-	-	South Korea, Russia, Singapore, Taiwan and GSO countries: Pre-harvest: 90d PHI as registered. Post-harvest as registered. Do not use in HOT water bath and not recommended for wax application.
Pyriproxyfen/ Nemesis	90d PHI as registered.	-	-	-	-	-	Switzerland: 120d PHI. Taiwan: 90d PHI as registered for oranges and soft citrus and not later than 90% petal fall for other citrus. South Korea: Not later than 90% petal fall.
Sodium ortho-phenylphenol (SOPP)	Post-harvest as registered.	-	-	-	-	-	South Korea: Not permitted.
Spinetoram/ Delegate	14d PHI for grapefruit and lemons and 7d PHI as registered of other citrus.	7d PHI as registered for oranges and soft citrus and 14d PHI for other citrus.	-	7d PHI as registered.	7d PHI as registered.	7d PHI as registered.	Great Britain ^m : 14d PHI.
Spinosad/ (Tracer/GF120)	1d PHI as registered for fruit fly baiting (GF120), but 28d PHI for other applications (Tracer).	-	-	-	-	-	-
Spirodiclofen/ Envidor	14d PHI as registered.	-	-	-	-	-	-
Spirotetramat/ Movento	60d PHI as registered.	-	-	-	-	-	-
Sulfoxaflor (Isoclast) / Closer	21d PHI as registered.	-	-	-	-	-	-
Sulphur	0d PHI as registered.	-	-	-	-	-	-
Tartar emetic/ Tartox	30d PHI as registered.	-	-	-	-	-	-
Tau-fluvalinate/ Klartan	Not later than mid-November as registered.	-	-	-	-	-	-
Tebuconazole/ Folicur	Not later than 90% petal fall as registered.	-	-	-	-	-	-
Temephos/ Abate	200d PHI	-	-	-	-	-	-

PRODUCT	All markets (including EU) except where other restrictions are specified	CODEX (A) ^a	CODEX (B) ^b	CANADA	U S A	JAPAN	Other ^d
Terbufos/ (AC92-100, Counter)	30d PHI as registered and not to be used between December and 1 April.	-	-	-	-	-	-
Tetradifon/ Tedion	Not later than 90% petal fall.	-	15d PHI	15d PHI	-	15d PHI	Switzerland: Not later than 90% petal fall. South Korea: 15d PHI for soft citrus and not later than 90% petal fall for other citrus.
Thiabendazole	Post-harvest as registered.	-	-	-	-	-	-
Thiacloprid/ Calypso	No later than 3 weeks after petal-fall.	-	-	-	-	-	-
Thiophanate-methyl/ Topsin	Not later than 90% petal fall.	-	14d PHI as registered.	14d PHI as registered.	Not later than 90% petal fall.	14d PHI as registered.	-
Trichlorfon/ Dipterex	28d PHI as registered.	-	10d PHI as registered.	-	-	-	-
<i>Trichoderma asperellum</i>	0d PHI as registered.	-	-	-	-	-	-
Trifloxystrobin/ Flint	76d PHI as registered.	-	-	-	-	-	Taiwan: Not later than 90% petal fall for soft citrus and 76d PHI as registered for other citrus.
Triflumuron/ Alsystin	Not later than 90% petal fall.	-	30d PHI as registered.	60d PHI	Not permitted	-	-
Uniconazole/ Tips Plus	135d PHI as registered.	-	-	-	-	-	-
2,4-D	Post-harvest: Not more than 250ppm in a packhouse treatment and not more than 250ppm in a pre-degreening drench. Pre-harvest:7d PHI as registered.	-	-	-	-	Pre-harvest as registered.	South Korea: Not permitted on oranges, soft citrus and grapefruit. China: °
3,5,6 TPA / Maxim	120d PHI as registered.	-	-	-	-	-	-

PHI = Pre-harvest interval.

- = as for "all markets", that is no additional restrictions apply.

GSO = Gulf State Organization (Bahrain, Kuwait, Oman, Qatar, Saudi Arabia, United Arab Emirates).

SUMMARY TABLE OF RESIDUE TOLERANCES

Chemical	RSA	General export tolerance	Codex	Canada	USA	Japan	South Korea
Acephate	0.2	0.01	None	0.1	None	0.01	5.0 ¹³ , 0.01 ¹¹
Acetamiprid	0.5	0.5	1.0	0.5	1.0	2.0	0.5
Acrinathrin	0.2	0.01	None	0.1	None	0.01	1.0 ¹³ , 0.01 ¹¹
Amitraz	0.2	0.05	0.5 ⁷ , None ¹¹	0.1	None	0.9 ^{7,13} , 0.5 ^{19,20}	0.7 ²⁰ , 0.2 ¹³ , 0.01 ^{7,19}
Avermectin	0.01	0.01	0.02	0.02	0.02	0.1	0.07 ¹³ , 0.05 ⁷ , 0.02 ²⁰ , 0.01 ¹⁹
Azadirachtin	None	0.01	None	0.1	Exempt	Exempt	0.01
Azinphos-methyl	2.0	0.01	1.0	2.0	None	2.0	0.01
Azoxystrobin	10.0	10.0	15.0	15.0	15.0	10.0	9.0 ¹³ , 10.0 ¹¹
<i>Bacillus thuringiensis</i>	None	None	None	None	None	None	Exempt
Bromopropylate	3.0	0.01	None	2.0	None	2.0	0.01
Buprofezin	0.05	0.01	1.0	4.0 ^{7,13} , 0.1 ^{19,20}	4.0	2.0 ⁷ , 3.0 ¹¹	2.5 ^{7,20} , 0.5 ¹³ , 0.01 ¹⁹
Cadusafos	0.05	0.01	None	0.1	None	0.01	0.01
Carbendazim	5.0	0.7 ^{13,20} , 0.2 ^{7,19} , (0.01 ^{7,19} , 0.9 ^{13,20}) ^{EU5}	1.0 ⁷ , None ¹¹	10.0	None	7.0 ^{19,20} , 3.0 ¹¹	5.0 ¹³ , 0.7 ²⁰ , 0.01 ¹¹
Chinomethionat	0.5	0.01	None	0.1	None	0.7	0.01
Chlorfenapyr	0.01	0.01	1.5 ⁷ , 0.8 ²⁰ , None ¹¹	0.1	0.01	2.0	1.0 ^{13,20} , 0.7 ⁷ , 0.01 ¹⁹
Chlorantraniliprole	0.5	0.5 (0.01) ^g	0.7	0.7	1.4	0.7	1.0 ^{13,20} , 0.6 ^{7,19}
Chlorpyrifos	0.3 (0.01) ^f	0.01	None	1.0	0.01	1.0	1.0
Clothianidin	0.01	0.01	0.07	0.1	0.07 ^{US1}	2.0	1.0 ¹³ , 0.3 ^{7,20} , 0.01 ¹⁹
Copper	20.0	20.0	None	50.0	Exempt	Exempt	Exempt
Cyantraniliprole	1.0	0.9	0.7	0.7	0.7	0.7	0.7
Cyhexatin (Azocytotin)	2.0	0.01, 0.2 ⁷	0.2 ⁷ , None ¹¹	0.1	Not permitted	0.5 ⁷ , 0.2 ¹¹	1.0 ¹³ , 0.01 ¹¹
Cypermethrin	0.2	2.0 (0.005 ⁷ , 0.3 ^{13,20} , 0.5 ¹⁹) ^{EU1}	0.3 ¹¹ , 0.5 ¹⁹	1.0	None, (0.35, 0.5 ¹⁹ , 0.3 ^{7,20}) ^c	2.0	2.0 ¹³ , 1.5 ²⁰ , 0.3 ^{7,19}
Dichlorprop	0.3	0.3	None	0.1	None	0.2	0.01
Dichlorvos	0.1	0.01	None	0.1	None	0.2	0.2 ¹³ , 0.05 ^{20,7} , 0.01 ¹⁹
Dicofol	5.0	0.02	None	0.1	None	5.0	0.01
Difenoconazole	0.05	0.05	0.6	0.8	0.6	0.6	0.6 ⁶ , 1.0 ¹³
Dimethoate	2.0	0.01	2.0 ¹³ , None ¹¹	1.5	2.0	2.0	0.01
Dimethyl Didecyl ammonium Chloride	6.0	0.05	None	0.1	Exempt	Not permitted ^k	0.01
Dipotassium phosphate	None	None	None	0.1	None	0.01	0.01
Dithiocarbamate	3.0	3.0	10.0 ¹³ , 2.0 ⁷ , None ¹¹	0.1	10.0 ¹³ , None ¹¹	10.0 ¹³ , 2.0 ¹¹	5.0 ¹³ , 3.0 ²⁰ , 0.01 ^{7,19}
(E)-8-Dodecen-1-yl acetate + (Z)-8-Dodecen-1-yl acetate	None	0.01	Exempt	None	Exempt	None	None
Emamectin Benzoate	0.01	0.002 ¹⁹ , 0.003 ¹¹	None	0.1	None	0.01	0.05 ^{7,13,20} , 0.01 ¹⁹
Ethephon	2.0	0.05	None	1.0	None	2.0	0.01

Chemical	RSA	General export tolerance	Codex	Canada	USA	Japan	South Korea
Ethoprophos	0.05	0.01	None	0.1	None	0.01	0.01
Etozazole	0.2	0.1 (0.01) EU4	0.1	0.1	0.1 ^{7,13} , None ¹¹	0.7	1.0 ¹³ , 0.7 ²⁰ , 0.01 ^{7,19}
Fenamiphos	0.05	0.01	None	0.1	None	0.04 ¹³ , 0.2 ¹¹ (0.01) ^{J1}	0.01
Fenazaquin	0.05	0.05	None	0.4	0.4	0.4	2.0 ¹³ , 0.4 ¹¹
Fenbutatin-oxide (Hexakis)	1.0	0.01	5.0	2.0	20.0	5.0	0.01
Fenpropathrin	0.5	0.5 (0.01) ^{EU2}	2.0	2.0	2.0	5.0	5.0 ¹³ , 2.0 ¹¹
Fenpyroximate	0.2	0.2	None ¹¹ , 1.0 ²⁰ , 0.5 ¹⁹	1.0	1.0	2.0	0.7
Fipronil	0.05	0.005	None	0.1	None	0.01	0.01
Fludioxonil	10.0	10.0	10.0	10.0	10.0	10.0	10.0
Fluopyram	0.01	0.01	1.0 ²⁰ , 0.6 ^{7,13} , 0.4 ¹⁹	1.0	1.0	1.0	1.0
Flutriafol	0.1	0.01	None	0.1	None	0.01	2.0 ^{13,20} , 0.01 ^{7,19}
Formetanate	0.5	0.01	None	0.9 ⁷ , 0.4 ¹⁹ , 0.09 ²⁰ , 0.03 ¹³	1.5 ^{7,19} , 0.6 ²⁰ , 0.03 ¹⁰	2.0 ^{7,19} , 0.6 ²⁰ , 0.03	0.01
Fosetyl-Al	50.0	50.0	50.0 ¹³ , 20.0 ⁷ , None ¹¹	9.0	9.0	150.0	9.0 ¹³ , 4.0 ⁷ , 0.01 ^{19,20}
Fosthiazate	0.1	0.02	None	0.1	None	0.01	0.01
Furfural	1.0 ⁿ	1.0	None	0.1	None	0.01	0.01
Gibberellic Acid	Exempt	Exempt	Exempt	0.1	Exempt	0.2	Exempt
Granulovirus	None	None	None	None	Exempt	None	None
Guazatine	5.0	0.05	5.0	0.1	None	0.01	0.01
<i>Helicoverpa armigera</i> nucleopolyhedrovirus	None	None	Exempt	None	None	None	None
Imazalil (Chloramizol)	5.0	5.0 ^{13,20} , 4.0 ^{7,19}	15.0	5.0	10.0	5.0	15.0
Imidacloprid	0.5	0.5 (0.01) ^{EU3}	1.0	1.0	0.7	0.7	0.7
Iprodione	1.0	0.01	None	0.07	None	10.0 (0.01) J2	7.0 ²⁰ , 2.0 ¹³ , 0.01 ^{7,19}
Isazophos	0.02	0.01	None	0.1	None	0.01	0.01
Kresoxim-methyl	0.5	0.5 ^{7,19} , 0.01 ¹¹	0.5 ^{7,19} , None ¹¹	9.0 ¹³ , 0.1 ¹¹	None	10.0	2.0 ¹³ , 0.01 ¹¹
Mercaptothion	4.0	2.0	7.0	0.1	8.0	7.0	4.0 ⁷ , 0.01 ¹¹
Metalaxyl-M (Mefenoxam)	1.0	0.5 ^{13,20} , 0.7 ^{7,19} (0.01) ²⁰ , 0.4 ¹³ , 0.7 ^{7,19} ^{EU6}	5.0	5.0	1.0	0.7	0.01
Methamidophos	0.2	0.01	None	0.1	None	0.01	0.2 ¹³ , 0.01 ¹¹
Methidathion	2.0	0.02	None	2.0	None	5.0	0.01
Methiocarb	0.1	0.03	None	0.1	None	0.05	0.5 ¹³ , 0.01 ¹¹
Methomyl (Thiodicarb)	0.2	0.01	1.0	1.0	2.0	10.0	0.7 ¹³ , 0.01 ¹¹
Methoxyfenozide	0.5	0.5	2.0	10.0	3.0	3.0	1.0 ¹³ , 3.0 ¹¹
Methyl-parathion	1.0	0.01	None	0.1	None	0.2	0.01
Mevinphos	0.1	0.01	None	0.2	None	0.01	0.01
Orange Oil	Exempt	Exempt	Exempt	None	Exempt	Exempt	Exempt
Omethoate	2.0	0.01	None	0.1	None	1.0	0.01
Parathion	0.5	0.05	None	0.1	None	0.5	0.01
Permethrin	0.01	0.01	0.5	0.1	None	5.0	0.01
Phenthoate	1.0	0.01	None	0.1	None	5.0	2.0 ²⁰ , 1.0 ¹¹

Chemical	RSA	General export tolerance	Codex	Canada	USA	Japan	South Korea
Phosphorous acid	50.0	50.0	None	0.1	Exempt	150.0	0.01
Pirimicarb	0.5	0.5	3.0	0.1	None	0.5 ⁷ , 0.05 ¹¹	0.01
Potassium bicarbonate	None	None	None	0.1	Exempt	0.01	0.01
Prochloraz	2.0	0.03	10.0	0.1	None	0.01	1.0 ¹³ , 0.01 ¹¹
Procymidone	0.2	0.01	None	0.1	None	5.0	0.01
Profenofos	1.0	0.01	None	0.1	None	0.01	0.01
Propiconazole	6.0	0.01	10.0, 4.0 ¹⁹	8.0	8.0	8.0	8.0
Prothiofos	0.05	0.01	None	0.1	None	0.01	0.01
Pymetrozine	0.3	0.02	None	0.2 ^{7,13,20} , 0.1 ¹⁹	None	0.01	0.3 ^{13,20} , 0.01 ^{7,19}
Pyraclostrobin	0.5	0.5	2.0	2.0	2.0	2.0	2.0
Pyrethrins (incl Pyrethrum)	1.0	1.0	0.05	1.0 ² , 0.1 ¹¹	1.0 ² , None ¹¹	1.0	0.01
Pyrimethanil	10.0	8.0	7.0	10.0	10.0	10.0	1.0 ¹³ , 7.0 ¹¹
Pyriproxyfen	0.2	0.2	0.5	0.5	0.5	2.0	0.7 ¹³ , 0.01 ¹¹
SOPP	10.0	10.0	10.0	10.0	10.0	10.0	0.01
Spinetoram	0.05	0.15 ¹³ , 0.07 ⁷ , 0.02 ^{19,20}	0.15 ¹³ , 0.07 ⁷ , None ¹¹	0.3	0.3	0.7	0.5 ¹³ , 0.05 ^{7,20} , 0.01 ¹⁹
Spinosad	0.05	0.05	0.3	0.3	0.3	0.3	0.3
Spirodiclofen	0.1	0.01	0.4	0.5	0.5	2.0	2.0 ¹³ , 0.4 ¹¹
Spirotetramat	1.0	0.5	0.5	0.6	0.6	3.0	2.0 ²⁰ , 0.5 ¹¹
Sulfoxaflor	0.3	0.3 ¹¹ , 0.15 ¹⁹	0.8 ^{7,13} , 0.4 ²⁰ , 0.15 ¹⁹	0.7	0.7	2.0	1.0 ¹³ , 0.7 ⁷ , 0.6 ²⁰ , 0.3 ¹⁹
Sulphur	50.0	50.0	Exempt	None	Exempt	Exempt	Exempt
Tartar emetic (Antimony)	3.0	1.0	None	0.1	None	0.01	0.01
Tau-Fluvalinate	1.0 ⁿ	0.4	None	0.1	None	2.0	0.01
Tebuconazole	0.02	0.02	0.7 ¹³ , 0.4 ⁷ , None ¹¹	1.0 ^{7,13} , 0.1 ¹¹	1.0 ⁷ , None ¹¹	5.0	2.0 ^{13,20} , 0.01 ^{7,19}
Temephos	1.0	0.01	None	0.1	None	0.01	0.01
Terbufos	0.1	0.01	None	0.1	None	0.005	0.01
Tetradifon	5.0	0.01	None	2.0	None	2.0	3.0 ¹³ , 0.01 ¹¹
Thiabendazole	6.0	6.0	7.0	10.0	10.0	10.0	10.0 ¹³ , 7.0 ¹¹
Thiacloprid	0.05	0.01	None	0.1	None	0.01	0.7 ^{7,20} , 0.3 ¹³ , 0.01 ¹⁹
Thiophanate-Methyl	5.0	0.01	None	10.0	None	7.0 ^{19,20} , 3.0 ¹¹	0.01
Trichlorfon	0.1	0.01	None	0.1	None	0.1	0.01
<i>Trichoderma asperellum</i>	None	None	Exempt	None	None	None	Exempt
Trifloxystrobin	0.1	0.1	0.5	0.6	0.6	3.0	0.5
Triflumuron	0.5	0.01	None	0.1	None	0.02	0.01
Uniconazole	0.01 ⁿ	0.01	None	0.1	None	0.01	0.01
2,4-D	2.0	1.0	1.0	2.0	3.0	3.0	1.0 ²⁰ , 0.15 ¹¹
3,5,6 TPA/Trichlopyr	0.1	0.1	None	0.1	None	0.1	0.1 ¹³ , 0.01 ¹¹

"None" = no MRL, therefore fruit must be free of detectable residue. However, in the case of CODEX countries additional options may apply – refer to notes.

SUMMARY TABLE OF CHANGES EFFECTIVE FROM THIS EDITION

<u>Active</u>	<u>Country</u>	<u>Previous MRL</u>	<u>New MRL</u>	<u>PHI Changes</u>
Carbendazim	EU	0.7 ^{13,20} , 0.2 ^{7,19}	0.7 ^{13,20} , 0.2 ^{7,19} (0.01 ^{7,19} , 0.9 ^{13,20}) ^{EU5}	120d PHI for oranges and grapefruit and 90d PHI for other citrus. (Not later than 90% petal fall for oranges and grapefruit and 90d PHI for lemons and mandarins) ^{EU5}
Chlorpyrifos	Codex	1.0	None	Not later than 90% petal fall for sprays, 0d PHI for soil and stem applications as registered.
Clothianidin	USA	N/A	N/A	The USA Clothianidin MRL expires on the 31st December 2026. (previously 2023).
Cypermethrin	EU	2.0	2.0 (0.005 ⁷ , 0.3 ^{13, 20} , 0.5 ¹⁹) ^{EU1}	28d PHI as registered. (Not later than 90% petal fall for oranges and 28d PHI as registered for other citrus). ^{EU1}
Dichlorvos	USA	N/A	N/A	21d PHI
Dithiocarbamates	GSO Countries	N/A	N/A	Not later than 90% petal fall for Lemons and Grapefruit and 21d or 28d PHI as registered for Soft Citrus and Oranges.
Dimethoate	Canada	N/A	N/A	14d PHI for foliar applications and 42d PHI for soil applications as registered.
Dimethoate	Codex	5.0	2.0 ¹³ , None ¹¹	14d PHI for foliar applications and 42d PHI for soil applications as registered for mandarins. For other citrus, not later than 50% petal fall for foliar applications and not later than white bud stage for soil applications.
Dimethoate	GSO	N/A	N/A	14d PHI for foliar applications and 42d PHI for soil applications as registered.
Dimethoate	Japan	N/A	N/A	14d PHI for foliar applications and 42d PHI for soil applications as registered.
Dimethoate	USA	N/A	N/A	14d PHI for foliar applications and 42d PHI for soil applications as registered.
Dipotassium phosphate	Canada	None	0.1	No change
Dipotassium phosphate	Japan	None	0.01	No change
Dipotassium phosphate	South Korea	None	0.01	No change
(E)-8-Dodecen-1-yl acetate + (Z)-8-Dodecen-1-yl acetate	Codex	None	Exempt	No change
(E)-8-Dodecen-1-yl acetate + (Z)-8-Dodecen-1-yl acetate	EU	None	Exempt	No change
Etoxazole	USA	0.1 ^{7, 13}	0.1 ^{7, 13} , None ¹¹	No change
Gibberellic Acid	Codex	None	Exempt	No change
<i>Helicoverpa armigera</i> nucleopolyhedrovirus	Codex	None	Exempt	No change
Metalaxyl-M	EU	0.7 ^{7,19} , 0.5 ^{13,20}	0.7 ^{7,19} , 0.5 ^{13,20} (0.01 ²⁰ , 0.4 ¹³ , 0.7 ^{7,19}) ^{EU6}	30d PHI as registered (Not later than 90% petal fall for lemons) ^{EU6}
Methidathion	Codex	5.0 ¹³ , None ¹¹	None	Not later than 90% petal fall.
Methidathion	South Korea	N/A	N/A	Not later than 90% petal fall.
Orange Oil	Codex	None	Exempt	No change
Orange oil	USA	None	Exempt	No change
Potassium bicarbonate	Canada	None	0.1	No change
Potassium bicarbonate	EU	None	Exempt	No change
Potassium bicarbonate	Japan	None	0.01	No change
Potassium bicarbonate	South Korea	None	0.01	No change
Potassium bicarbonate	USA	None	Exempt	No change
Pyrethrins	Canada	1.0 ²	1.0 ² , 0.1 ¹¹	No change

Pyrethrins	GSO Countries	N/A	N/A	Not later than 90% petal fall.
Pyrethrins	USA	1.0 ²	1.0 ² , None ¹¹	No change
Pyrimethanil	GSO Countries	N/A	N/A	Pre-harvest: 90d PHI as registered. Post-harvest as registered. Do not use in HOT water bath and not recommended for wax application.
Pyriproxyfen	EU	0.6	0.7	No change
Sulphur	Codex	None	Exempt	No change

NOTES

Numerical Superscripts:

- 1 fruit without peel/pulp
- 2 just oranges
- 3 exocarp of summer oranges
- 4 fruit (except exocarp of summer oranges)
- 5 just peel
- 6 except mandarin oranges
- 7 Oranges, sweet, sour
- 8 Citrus pulp, dried
- 9 Whole fruit
- 10 Clementines, soft citrus
- 11 Other citrus
- 12 Except summer oranges
- 13 Soft citrus
- 14 Pulp juice
- 15 No specific crop
- 16 Summer orange and soft citrus
- 17 Summer orange, pulp and peel
- 18 Summer orange, pulp
- 19 Only grapefruit
- 20 Only lemons
- 21 Soft citrus, limes and lemons
- 22 Grapefruit, oranges and pommelos.

“None” = no MRL, therefore fruit must be free of detectable residue. However, in the case of CODEX countries additional options may apply – refer to notes.

Alphabetical Superscripts:

a = CODEX (A): See Regions and Country Groupings Table below.

b = CODEX (B): See Regions and Country Groupings Table below.

c = USA has MRLs on citrus for Alpha-Cypermethrin (0.35 mg/kg) and Zeta-Cypermethrin (0.5 mg/kg for grapefruit, 0.3 mg/kg for lemons and oranges), but none for Cypermethrin, which means only Alpha and Zeta-Cypermethrin formulations are suitable for the USA.

d = Reference to “Citrus Fruits” in Taiwan MRL legislation should be interpreted as “oranges only” but “Citrus” refers to all citrus types.

EU1 = The EU notified their intention to lower the Cypermethrin MRL. If adopted a restriction of not later than 90% petal fall will apply for oranges and 28d PHI as registered for other citrus with the expected date of entry into force in October, 2024. (WTO notification: G/SPS/N/EU/702).

EU2 = The EU Fenpropathrin MRL is expected to be lowered to LOD no earlier than May, 2023. A 180d PHI will then apply.

EU3 = EU Imidacloprid MRLs are under review and will be lowered with the expectation that this will occur in the second half of 2022 or in the first quarter of 2023, subject to the EU MRL decision-making process.

EU4 = The EU have notified of their intention to lower the Etoxazole MRL to LOD with the proposed date of entering into force 1st of May, 2024 (WTO notification: G/SPS/N/EU/617).

EU5 = The EU have notified of their intention to lower the Carbendazim MRLs for grapefruit and oranges to 0.01 ppm and to set new Carbendazim MRLs for lemons and mandarins at 0.9 ppm. The new MRLs for lemons and mandarins are expected to apply from January 2025. All oranges and grapefruit in the EU market from July, 2024 should comply with the lower MRL of 0.01 ppm. (WTO notification: G/SPS/N/EU/696).

EU6 = The EU notified their intention to lower the Metalaxyl-M MRL to 0.01 ppm for lemons and 0.4 ppm for mandarins with the proposed date of entering into force December, 2024. If adopted, a restriction of not later than 90% petal fall will then apply to lemons (WTO notification: G/SPS/N/EU/697).

f = The Registrar has indicated the imminent phase out of Chlorpyrifos registrations and the MRL will be lowered to LOD (WTO notification: G/SPS/N/ZAF/81, Gazette NO. 48847 of June 2023).

g = Norway has a new Chlorantraniliprole (Rynaxypyr) MRL of 0.01mg/kg. A PHI of not later than 90% petal fall applies.

J1 = Japan has notified their intention to lower the Fenamiphos MRL to LOD (WTO notification G/SPS/N/JPN/1216).

J2 = Japan has notified their intention to lower the Iprodione MRL to LOD (WTO G/SPS/N/JPN/1218).

k = Pending decision by Japanese authorities regarding the use of disinfectants.

l = See CRI Production Guideline for appropriate application techniques.

m = Great Britain includes England, Scotland and Wales. EU MRLs apply in Ireland.

n = This active has obtained RSA usage authorization and a provisional MRL. CGA is engaged in a process with the Registrar and Department of Health to confirm these MRLs.

o = China set the 2,4-D MRL to 0.1 mg/kg for oranges and soft citrus in February 2020. Packhouse managers need to determine practices in their situation which will result in residues below 0.1 mg/kg for oranges and soft citrus.

q = Gulf states that subscribe to the Gulf Standards Organization (GSO) MRL include UAE, Kingdom of Bahrain, Kingdom of Saudi Arabia, Sultanate of Oman, State of Kuwait and Republic of Yemen. The GSO has published a 2,4-D MRL of 1.0 mg/kg.

T1 = Taiwan has a Grapefruit MRL of 5.0 mg/kg and use of Fludioxonil as registered may lead to exceedances of this MRL.

US1 = The USA Clothianidin MRL expires on the 31st December 2026.

Y = Pay special attention to the introductory notes on page one (3rd paragraph) dealing with lemons

REGIONS AND COUNTRY GROUPINGS

<u>Description</u>	<u>Countries</u>
CODEX A (Where no national MRL is set, CODEX MRLs apply)	Africa [Angola, Benin, Botswana, Congo (Republic of), Gabon, Kenya, Madagascar, Mali, Mauritius*, Mauritania, Namibia, Senegal, Seychelles, Reunion*, Sudan*, Tanzania] Asia [China (Peoples Republic of), Hong Kong, Indonesia, Malaysia, Singapore, Thailand, Vietnam]
CODEX B (Where no national MRL is set, CODEX MRLs apply, or when no CODEX MRL is set RSA MRL apply)	Africa [Burkina Faso, Cameroon, Cote D'ivoire, Malawi, Nigeria, Tunisia, Uganda] Asia [Bangladesh, Sri Lanka, Philippines] Middle East [Azerbaijan, Iran, Jordan*, Pakistan] Other [Russian Federation, Georgia*]
Gulf State Organization (GSO)	Bahrain, Kuwait, Oman, Qatar, Saudi Arabia, United Arab Emirates

* It has not been possible to re-confirm these requirements recently but presumably they still apply